## **Return to Work Guidance After COVID-19 Illness or Exposure**

	T	
Symptomatic with confirmed	Mild or moderate illness and not	Severe to critical illness or
COVID-19 or suspected COVID-19	severely immunocompromised:	severely immunocompromised*:
	Return to work when: Minimum	Return to work when: Minimum
	of 10 days have passed since	of 20 days have passed since
	symptoms <u>first appeared</u> , AND	symptoms <u>first appeared</u> , AND
	minimum of 24 hours have passed	minimum of 24 hours have passed
	since last fever without	since last fever without
	medications, AND symptoms have	medications, AND symptoms have
	improved	improved
Asymptomatic with confirmed	Not severely	Severely immunocompromised*:
COVID-19	immunocompromised:	
		Return to work when: Minimum
	Return to work when: Minimum	of 20 days have passed since
	of 10 days have passed since	positive lab test AND remains
	positive lab test AND remains	asymptomatic
	asymptomatic	
Asymptomatic with known	Return to work and follow these	
exposure to COVID-19 working in	practices:	
critical infrastructure		
	Employer prescreens employee	
	prior to employee's building entry,	
	AND employee monitors	
	temperature and symptoms, AND	
	employee wears mask for 14 days	
	after last exposure, AND	
	employee practices social	
	distancing while at work (as able),	
	AND both employee and employer	
	routinely disinfect and clean	
	impacted work spaces	
	If the employee becomes sick	
	during the day, the employer will:	
	send the employee home, clean	
	and disinfect work spaces, and	
	compile a list of contacts as	
	outlined in CDC's Interim	
	Guidance Protocol.	

<sup>\*</sup> NOTE: 'The degree of immunocompromised for the employee is determined by the treating provider.' If severely immunocompromised, documentation stating such must be generated by the treating provider and given to the employer.